

Josephson current through a Luttinger liquid

Rosario Fazio⁽¹⁾, F. W. J. Hekking⁽²⁾, and A. A. Odintsov^(2,3)

⁽¹⁾*Istituto di Fisica, Università di Catania, viale A. Doria 6, 95129 Catania, Italy*

⁽²⁾*Institut für Theoretische Festkörperphysik, Universität Karlsruhe, Postfach 6980, 76128
Karlsruhe, FRG*

⁽³⁾*Present address: Electrotechnical Laboratory, 1-1-4 Umezono, Tsukuba-shi, Ibaraki 305, Japan*
*Permanent address: Nuclear Physics Institute, Moscow State University, Moscow 119899 GSP,
Russia*

(February 1, 2008)

Abstract

We study the Josephson effect through a one-dimensional system of interacting electrons, connected to two superconductors by tunnel junctions. The interactions are treated in the framework of the one-channel Luttinger model. At zero temperature, the Josephson critical current is found to decay algebraically with increasing distance between the junctions. The exponent is proportional to the strength of the Coulomb interaction. If the Luttinger liquid has a finite size, the Josephson current depends on the total number of electrons modulo 4. These parity effects are studied for the ring, coupled capacitively to a gate-voltage and threaded by a magnetic flux. The Josephson current changes continuously as a function of the gate voltage and *stepwise* as a function of the magnetic flux. The electron-electron interaction introduces *qualitatively* new features compared to the non-interacting case.

PACS numbers: 74.50 +r, 72.15 Nj

Recent technological developments in the fabrication of semiconductor- superconductor (S-Sc) interfaces made the observation of supercurrent in S-Sc-S junctions possible¹. This progress was accompanied by a number of appealing theoretical predictions. The Josephson current through a narrow constriction in a high-mobility two-dimensional non-interacting electron gas (2DEG) should be quantized². Interference effects of two electrons entering the normal region from a superconductor are important, especially in mesoscopic samples³. This interference influences the Josephson current through the normal region.

It is well-known that electron-electron interactions affect the transport properties of mesoscopic devices as well. The interaction manifests itself in the single charge effects⁴. In particular, the Josephson current through SSS and SNS systems can be modulated by a gate-voltage⁵. This has been described in terms of a phenomenological capacitance model. However, in low-dimensional semiconductor nanostructures, with a low electron density, electron-electron interactions should be treated microscopically. As a result, the transport properties of a (quasi-) one-dimensional quantum wire show deviations from Fermi-liquid behaviour⁶.

In the present paper we analyze the Josephson current through a Luttinger liquid (LL). Specifically, we consider two geometries which can be realized experimentally: A long wire with tunnel contacts to two superconductors at a distance d (Fig. 1a), and a ring with circumference L and tunnel contacts at a distance $L/2$ (Fig. 1b). Various aspects of transport in mesoscopic systems (parity effects and interference combined with electron-electron interactions) and their interplay can be studied in this system. The aim of this paper is twofold. First, we want to study the influence of the Coulomb interaction on the Josephson critical current. Second, we would like to see how the parity effects which are present in the ring^{9–12} manifest themselves in the dependence of the Josephson current on magnetic flux and gate-voltage. The coupling of bulk superconductors to a one-dimensional (1D) system of interacting electrons was studied by Fisher⁷. In contrast to our work, a chiral LL of spin-polarized electrons was considered and tunneling with spin flip was analyzed.

The systems under consideration (Fig. 1) can be described by the Hamiltonian $\hat{H} = \hat{H}_{S1} + \hat{H}_{S2} + \hat{H}_L + \hat{H}_T$. Here, $\hat{H}_{S1}$, $\hat{H}_{S2}$ are the BCS-Hamiltonians for the bulk superconductors with gap Δ_1 , Δ_2 respectively, kept at a phase difference $\chi = \chi_1 - \chi_2$. The (1D) electron system is described by the Hamiltonian⁸

$$\hat{H}_L = \hbar \int \frac{dx}{\pi} \sum_j v_j \left[\frac{g_j}{2} (\nabla \phi_j)^2 + \frac{2}{g_j} (\nabla \theta_j)^2 \right]. \quad (1)$$

It is written as a sum of the contributions from the spin ($j = \sigma$) and charge ($j = \rho$) degrees of freedom. We use the standard notation⁸ for the parameters g_j of the interaction strengths and the velocities v_j of spin and charge excitations. We also introduce the bosonic fields $\phi_s = \phi_\rho + s\phi_\sigma$ and $\theta_s = \theta_\rho + s\theta_\sigma$ for spin up ($s = 1$) and down ($s = -1$) fermions. These fields obey the commutation relation $[\phi_s(x), \theta_{s'}(x')] = (i\pi/2) \text{sign}(x' - x) \delta_{s,s'}$. The tunneling is assumed to occur through two tunnel junctions at the points $x = 0$ and $x = d$,

$$\hat{H}_T = \sum_s T_1 \hat{\Psi}_{S1,s}^\dagger(x=0) \hat{\Psi}_{L,s}(x=0) + T_2 \hat{\Psi}_{S2,s}^\dagger(x=d) \hat{\Psi}_{L,s}(x=d) + (\text{h.c.}). \quad (2)$$

The tunnel matrix elements $T_{1,2}$ can be related to the tunnel conductances $G_{1,2}$ of the junctions: $G_i = (4\pi e^2/\hbar) N_L(0) N_i(0) T_i^2$, where $N_L(0) = 1/\pi\hbar v_F$, N_i is the normal density of states of a superconductor ($i = 1, 2$), and v_F is the Fermi velocity.

The fermionic field operators $\hat{\Psi}$ can be expressed in terms of spin and charge degrees of freedom⁹:

$$\hat{\Psi}_{L,s}^\dagger(x, \tau) = \sqrt{\rho_{0,s}} \sum_{m, \text{odd}} \exp \{imk_F x\} \exp \{im\theta_s\} \exp \{i\phi_s\}, \quad (3)$$

where k_F is the Fermi-wave vector and $\rho_{0,s} \equiv N_s/L = k_F/(2\pi)$ is the average electron density for one spin direction. If the LL is confined to a ring¹⁰, threaded by an Aharonov-Bohm flux Φ , the twisted boundary condition $\hat{\Psi}_{L,s}^\dagger(x+L) = \exp \{-2\pi i\Phi/\Phi_0\} \hat{\Psi}_{L,s}^\dagger(x)$ can be imposed¹¹. Here, Φ_0 is the normal flux quantum h/e . The fields θ and ϕ are decomposed in terms of bosonic fields and topological excitations⁹ in the following fashion:

$$\begin{aligned} \theta_j(x) &= \bar{\theta}_j(x) + \theta_j^0 + \pi M_j(x/2L), \\ \phi_j(x) &= \bar{\phi}_j(x) + \phi_j^0 + \pi(J_j - 4\delta_{j,\rho}\Phi/\Phi_0)(x/2L). \end{aligned} \quad (4)$$

Here, $\bar{\theta}_j$ and $\bar{\phi}_j$ are the non-zero modes

$$\begin{aligned} \bar{\theta}_j(x) &= \frac{i}{2} \sqrt{\frac{g_j}{2}} \sum_{q \neq 0} \left| \frac{\pi}{qL} \right|^{1/2} \text{sign}(q) e^{iqx} (\hat{b}_{j,q}^\dagger + \hat{b}_{j,-q}), \\ \bar{\phi}_j(x) &= \frac{i}{2} \sqrt{\frac{2}{g_j}} \sum_{q \neq 0} \left| \frac{\pi}{qL} \right|^{1/2} e^{iqx} (\hat{b}_{j,q}^\dagger - \hat{b}_{j,-q}), \end{aligned} \quad (5)$$

where $\hat{b}_{j,q}, \hat{b}_{j,q}^\dagger$ are Bose operators. M_j and J_j denote the topological excitations. They are related to the usual topological excitations for fermions with spin s : $M_s = (1/2)[M_\rho + sM_\sigma]$ and $J_s = (1/2)[J_\rho + sJ_\sigma]$. Using the topological constraints for M_s and J_s given in Ref.⁹, we find constraints for M_j and J_j : (i) The topological numbers M_j and J_j are either *simultaneously* even or *simultaneously* odd; (ii) when N_s is *odd* the sum $M_\rho \pm M_\sigma + J_\rho \pm J_\sigma$ takes values $\dots, -4, 0, 4, \dots$, when N_s is *even* the sum $M_\rho \pm M_\sigma + J_\rho \pm J_\sigma$ takes values $\dots, -6, -2, 2, 6, \dots$. Here, the number of electrons N_s ($N_{s=1} = N_{s=-1}$) determines the linearization point $k_F \equiv \pi N_s/L$ of the original electron spectrum¹⁰.

The Hamiltonian is decoupled in the non-zero modes and the topological excitations:

$$\hat{H}_L = \hbar \sum_{j=\rho, \sigma} \left\{ \sum_{q \neq 0} v_j |q| \hat{b}_{q,j}^\dagger \hat{b}_{q,j} + \frac{\pi v_j}{4L} \left[\frac{g_j}{2} (J_j - 4\delta_{j,\rho} f_\Phi)^2 + \frac{2}{g_j} (M_j - 4\delta_{j,\rho} f_\mu)^2 \right] \right\}. \quad (6)$$

We introduced the flux frustration $f_\Phi \equiv \Phi/\Phi_0$, as well as the parameter $f_\mu = (g_\rho L/4\pi v_\rho) \Delta\mu$, which is related to the difference $\Delta\mu$ between the electro-chemical potential μ of the superconducting electrodes and the Fermi energy $E_{F,0}$ of the ring. For non-interacting electrons the latter is just the energy $(\hbar k_F)^2/2M$ in the linearization point. Generally, the reference point $\Delta\mu = 0$ is defined from the requirements that for $\Phi = 0$ there are $2N_s$ electrons in the ground state ($N_1 = N_{-1}$) and the energies to add and remove the electron(s) to/from the system are the same. The electro-chemical potential μ can be controlled by a gate-voltage.

The stationary Josephson effect can be obtained by evaluating the phase-dependent part of the free energy $\mathcal{F}(\chi)$: The Josephson current is given by $I_J = -(2e/\hbar) \partial \mathcal{F} / \partial \chi$. We expand $\mathcal{F} = -(1/\beta) \ln Z$ where $Z = \text{Tr} \exp \{-\beta \hat{H}\}$ in powers of $\hat{H}_T$; the lowest order contribution arises in 4th order. It is represented by the diagram in the inset of Fig. 2, which

has a clear physical meaning: the Josephson effect occurs through a process which transfers a Cooper pair from superconductor $S2$, described by the anomalous Green's function $F_{S2}(\tau_3, \tau_4)$ through the second junction with an amplitude $(T_2^*)^2$. Subsequently, the two electrons propagate as a Cooperon through the LL from contact 2 to 1. This propagation is determined by the Cooperon propagator $\Pi(0, d; \tau_1, \dots, \tau_4)$. Finally, both electrons tunnel through the first junction (amplitude T_1^2) and enter the superconductor $S1$ as a Cooper pair, characterized by the anomalous function $F_{S1}^\dagger(\tau_1, \tau_2)$. If the distance d between the junctions is much larger than the coherence length in the superconductor, the characteristic energies $\hbar v_F/d$ of the electrons propagating through the 1D system are much less than the energies Δ of excitations in superconductors. At low temperatures ($k_B T \ll \Delta$), a generic process consists of fast tunneling of two electrons from the superconductor into the 1D system ($|\tau_1 - \tau_2| \sim \hbar/\Delta$) and their slow propagation through the LL ($|\tau_1 - \tau_3| \sim d/v_F$). The phase-dependent part of the free energy then simplifies to

$$\mathcal{F}(\chi) = -2\pi^2 N_1(0) N_2(0) \Re[T_1^2 (T_2^*)^2 e^{-i\chi} \int_0^\beta d\tau \Pi(d; \tau)], \quad (7)$$

with the Cooperon

$$\Pi(d, \tau) = \langle T_\tau \hat{\Psi}_{L,+}(0, 0) \hat{\Psi}_{L,-}(0, 0) \hat{\Psi}_{L,-}^\dagger(d, \tau) \hat{\Psi}_{L,+}^\dagger(d, \tau) \rangle, \quad (8)$$

where the average is taken over the eigenstates of $\hat{H}_L$. The evaluation of (8) with the help of bosonized field operators like (3) is straightforward.

Wire geometry. For an infinitely long wire (Fig. 1a), the topological excitations play no role since their energies $\sim \pi v_j/L$ are vanishingly small. The Cooperon propagator $\Pi(d, \tau)$ is given by

$$\Pi_w(d, \tau) = \rho_0^2 \sum_{n_1, n_2, \text{odd}} e^{i(n_1+n_2)(k_F d + \eta)} \times \left[\frac{\alpha^2}{d^2 + v_\sigma^2 \tau^2} \right]^{g_\sigma(n_1-n_2)^2/16} \left[\frac{\alpha^2}{d^2 + v_\rho^2 \tau^2} \right]^{1/g_\rho + g_\rho(n_1+n_2)^2/16}, \quad (9)$$

where $e^{i\eta} = [(d + i v_\rho \tau)/(d - i v_\rho \tau)]^{1/2}$, and α is a cut-off parameter¹³ of the order $1/k_F$. If $n_1 + n_2 \neq 0$, the rapid oscillations related to $e^{i(n_1+n_2)(k_F d)}$ make the Josephson term vanishingly small if the tunnel junctions are large compared to the Fermi wavelength. For $n_1 + n_2 = 0$ the leading contributions to the correlation function correspond to $n_1 - n_2 = \pm 2$. For the spin-independent electron-electron interaction we should fix $g_\sigma = 2$ and $v_\rho = (2/g_\rho)v_F$. For the Josephson current at zero temperature we find

$$I_J = \frac{2\pi e v_F}{d} \frac{G_1 G_2}{(4e^2/\hbar)^2} F_w(g_\rho, d) \sin(\chi), \quad (10)$$

$$F_w = \left[\frac{1}{k_F d} \right]^{2/g_\rho - 1} \int \frac{dx}{\pi} \frac{1}{\sqrt{1+x^2}} \left[\frac{1}{1 + (2x/g_\rho)^2} \right]^{1/g_\rho} \quad (11)$$

($F_w = 1$ for noninteracting electrons).

Ring geometry. For the ring (Fig. 1b) we get

$$I_J = \frac{2\pi e v_F}{L} \frac{G_1 G_2}{(4e^2/\hbar)^2} \sum_{\epsilon=\pm 1} \langle F_r(g_\rho, L, \epsilon, M_\sigma, J_\rho) \sin(\chi + \epsilon\pi M_\sigma/2 + \pi J_\rho/2) \rangle_{J,M}, \quad (12)$$

with

$$F_r = \left[\frac{\pi}{k_F L} \right]^{2/g_\rho-1} \int dx \frac{1}{\cosh(x)} \left[\frac{1}{\cosh(2x/g_\rho)} \right]^{2/g_\rho} \cosh \left[\left(\frac{2}{g_\rho} \right)^2 (M_\rho - 4f_\mu)x + \epsilon J_\sigma x \right]. \quad (13)$$

Here, $\langle \dots \rangle_{J,M}$ means evaluation with respect to the ground-state configuration of the topological excitations J_j , M_j of the LL subject to the topological constraints.

In Fig. 2 the dependence of the critical current on g_ρ for the two geometries is shown. The Josephson current is suppressed by the repulsive interaction, $I_J \propto d^{-2/g_\rho}$. We estimate the magnitude of the current, using typical numbers from the experiment of Mailly et al.¹⁴, to be of the order of several nA in the non-interacting case.

We discuss now the flux and gate-voltage dependence in the ring geometry. The Josephson current Eq. (13) depends on the gate-voltage via the parameter f_μ . The flux does not enter explicitly into Eq. (13). However, the Josephson current depends on f_Φ (and on f_μ) *implicitly* via the topological numbers $(J_\rho, J_\sigma, M_\rho, M_\sigma)$. Therefore, one can expect that the Josephson current changes stepwise as a function of the flux (the jumps correspond to the change in the topological numbers) and shows a continuous dependence (with jumps) on the gate-voltage (see Figs. 3, 4).

Without loss of generality we restrict the further consideration by odd values of N_s ¹⁵. We start from the non-interacting case $g_\rho = 2$. The ground-state configurations $(J_\rho, J_\sigma, M_\rho, M_\sigma)$ are displayed in Fig. 3a. Each state in the ring below the chemical potential μ of the superconductors is occupied by two electrons with opposite spin. Hence, the topological numbers M_σ and J_σ are always zero. The increase of the flux shifts down (up) the energy levels of the electrons moving in the positive (negative) direction along the ring. Increasing f_μ corresponds to a uniform shift of all energy levels down with respect to the chemical potential μ . A *pair* of electrons tunnels into (out of) the ring each time when an empty (filled) energy level crosses the chemical potential. This leads to a change in the topological number M_ρ , $\Delta M_\rho = 2(-2)$. Simultaneously, the topological number J_ρ changes, $\Delta J_\rho = \pm 2$, since the system acquires orbital momentum. With the increase of the flux Φ by one flux quantum, a pair of electrons enters the positive branch of the spectrum and (for a different value of Φ) another pair leaves the negative branch. For each process $\Delta J_\rho = 2$. Accordingly, the Josephson critical current $I_{J,c} \equiv \max_\chi |I_J(\chi)|$ (Fig. 3) changes stepwise with the flux making two jumps per period Φ_0 . One of the jumps occurs at $f_\Phi = -f_\mu + 1/2$ in Fig. 3.

For $f_\mu = 0$, the chemical potential of the superconductors μ is in the middle of the gap between the last occupied and first unoccupied energy levels ($E_{F,0} = \mu$). For this reason, the two processes mentioned above occur at the same point $f_\sigma = 1/2$. One can say that a pair of electrons jumps from the negative branch to positive branch (e.g. the transition $(0, 0, 0, 0) \rightarrow (4, 0, 0, 0)$ in Fig. 3a) and the number of electrons in the system is conserved.

The situation changes drastically for interacting electrons. The interaction lifts the degeneracy of the single particle states with different spin and the system acquires additional stiffness with respect to the change in the electron number (for the repulsive interactions $g_\rho < 2$ the coefficient $2/g_\rho$ of the term with the particle number M_ρ is larger than the coefficients of the other topological terms in Eq. (6)). For this reason, the electrons tunnel *one by one*

into the ring and new ground-state configurations (e.g. $(J_\rho, J_\sigma, M_\rho, M_\sigma) = (1, 1, 1, 1)$ and $(3, -1, 11)$) with an *odd* number of electrons arise (see Fig. 4a). In particular, the parity of the electron number on the ring can be changed not only by changing the gate-voltage, but also by changing the flux. Depending on the value of the gate-voltage the increase of the flux by one flux quantum is accompanied by the tunneling of two, one or zero electrons into the ring and the same number of tunneling events out of it. Accordingly, the Josephson current (Fig. 4b) shows four, two or zero jumps per period Φ_0 .

The size of the intervals of the gate-voltage, where the electron number (characterized by M_ρ) does not depend on the flux, increases with the increase of the repulsive interaction. In the limit of strong interaction the number of particles is fixed for any value of the gate-voltage except for small intervals near the points $f_\mu = 1/8 + n/4$ with integer n ¹⁶.

We have also found that for some ground-state configurations, the Josephson current changes sign and the ring acts as a π -junction. This happens, for example, in the regions with $J_\rho = 2 \bmod(4)$ both in the interacting and in the non-interacting cases (e.g. in the region $(2,0,2,0)$ in Figs. 3a, 4a). Such behaviour can be tested through the interference pattern of a SQUID, consisting of the ring and a standard Josephson junction. A more detailed analysis of all these features will be presented elsewhere¹⁷.

In this letter we studied the Josephson effect through a 1D system of interacting electrons: a quantum wire and a ring. The Josephson current was found to be suppressed by the Coulomb interaction. The interaction qualitatively modifies the parity effects in the ring and effects the dependences of the critical current on the flux and gate-voltage. An anomalous sign of the Josephson current was found in some range of parameters.

Acknowledgements We would like to thank L. Glazman, D. Khmel'nitskii, A.I. Larkin, D. Loss, and G.T. Zimanyi for useful discussions. The financial support of the European Community (HCM-network CHRX-CT93-0136) and of the Deutsche Forschungsgemeinschaft through SFB 195 is gratefully acknowledged.

REFERENCES

- ¹ J. Nitta et al., Phys. Rev. B **46**, 14286 (1992); A. Dimoulas, J.P. Heida, B.J. van Wees, T.M. Klapwijk, W. v.d. Graaf, and G. Borghs (unpublished)
- ² C.W.J. Beenakker and H. van Houten, Phys. Rev. Lett. **66**, 3056 (1991); A. Furusaki et al., ibid. **67**, 132 (1991)
- ³ B.J. van Wees et al., Phys. Rev. Lett. **69**, 510 (1992); F.W.J. Hekking and Yu.V. Nazarov, Phys. Rev. Lett. **71**, 1625 (1993)
- ⁴ D.V. Averin and K.K. Likharev, in “*Mesoscopic Phenomena in Solids*”, edited by B.L. Altshuler, P.A. Lee, and R. Webb (North- Holland, Amsterdam, 1991); G. Schön and A.D. Zaikin, Phys. Rep. **198**, 237 (1990)
- ⁵ K.A. Matveev et al., Phys. Rev. Lett. **70**, 2940 (1993); P. Joyez et al., ibid. **72**, 2458 (1994); R. Bauernschmitt et al., Phys. Rev. B **49**, 4076 (1994)
- ⁶ C.L. Kane and M.P.A. Fisher, Phys. Rev. Lett. **68**, 1220 (1992); K.A. Matveev and L.I. Glazman, ibid. **70**, 990 (1993)
- ⁷ M.P.A. Fisher (unpublished)
- ⁸ C.L. Kane and M.P.A. Fisher, Phys. Rev. B **46**, 15233 (1992)
- ⁹ F.D.M. Haldane, Phys. Rev. Lett. **47**, 1840 (1981); J. Phys. C **14**, 2585 (1981)
- ¹⁰ D. Loss, Phys. Rev. Lett. **69**, 343 (1992)
- ¹¹ N. Byers and C.N. Yang, Phys. Rev. Lett. **7**, 46 (1961). Note that the corresponding gauge transformation leads to an additional phase factor for the tunnel matrix element T_2 : $T_2 \rightarrow T_2 \exp \{-i\pi\Phi/\Phi_0\}$ for the set-up depicted in Fig. 1b.
- ¹² S. Fujimoto and N. Kawakami (unpublished)
- ¹³ A. Luther and I. Peschel, Phys. Rev. B **9**, 2911 (1974)
- ¹⁴ D. Mailly et al. Phys. Rev. Lett. **70**, 2020 (1993)
- ¹⁵ The case of even N_s leads to a similar picture apart from a relative shift along the f_Φ and f_μ axes.
- ¹⁶ This regime resembles the situation in small metallic tunnel junctions where the charging energy is much larger than the electron level spacing.
- ¹⁷ R. Fazio, F.W.J. Hekking, and A.A. Odintsov (unpublished)

FIGURES

FIG. 1. The geometries discussed in the text: (a) one-dimensional wire connected to two superconductors by tunnel junctions, separated by a distance d . (b) Ring with circumference L , threaded by a magnetic flux Φ . It is connected to two superconductors by tunnel junctions, separated by a distance $L/2$.

FIG. 2. Critical current as a function of g_ρ for the ring (solid line, normalized to $I_{J,c}^{(0)} = (4\pi ev_F/L)(G_1 G_2/(4e^2/\hbar)^2)$) and the wire (dashed line, normalized to $I_{J,c}^{(0)} = (\pi ev_F/d)(G_1 G_2/(4e^2/\hbar)^2)$). Inset: Lowest-order contribution to the Josephson effect.

FIG. 3. (a) Ground-state configurations for topological quantum numbers $(J_\rho, J_\sigma, M_\rho, M_\sigma)$ as a function of gate-voltage (f_μ) and magnetic flux (f_Φ), for $g_\rho = 2$. (b) Critical current at $T = 0$ for $g_\rho = 2$ as a function of f_Φ and f_μ .

FIG. 4. The same as Fig. 3, but for $g_\rho = 1.75$.